

# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Decreasing PICU Catheter-Associated Bloodstream Infections: NACHRI's Quality Transformation Efforts**

Marlene R. Miller, Michael Griswold, J. Mitchell Harris, II, Gayane Yenokyan, W. Charles Huskins, Michele Moss, Tom B. Rice, Debra Ridling, Deborah Campbell, Peter Margolis, Stephen Muething and Richard J. Brill

*Pediatrics* published online Jan 11, 2010;

DOI: 10.1542/peds.2009-1382

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.pediatrics.org>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# Decreasing PICU Catheter-Associated Bloodstream Infections: NACHRI's Quality Transformation Efforts



**WHAT'S KNOWN ON THIS SUBJECT:** CA-BSIs are a significant source of morbidity, mortality, and added medical costs to hospitalized adult and pediatric patients. Despite these data, strategies for reducing health care complications such as pediatric CA-BSIs have received relatively little attention.



**WHAT THIS STUDY ADDS:** In contrast with adult ICU care, maximizing insertion-bundle compliance alone cannot help PICUs to eliminate CA-BSIs. Instead, the main drivers for additional reductions in pediatric CA-BSI rates seem to be issues surrounding daily maintenance care for central lines.

## abstract

FREE

**OBJECTIVE:** Despite the magnitude of the problem of catheter-associated bloodstream infections (CA-BSIs) in children, relatively little research has been performed to identify effective strategies to reduce these complications. In this study, we aimed to develop and evaluate effective catheter-care practices to reduce pediatric CA-BSIs.

**STUDY DESIGN AND METHODS:** Our study was a multi-institutional, interrupted time-series design with historical control data and was conducted in 29 PICUs across the United States. Two central venous catheter-care practice bundles comprised our intervention: the insertion bundle of pediatric-tailored care elements derived from adult efforts and the maintenance bundle derived from the Centers for Disease Control and Prevention recommendations and expert pediatric clinician consensus. The bundles were deployed with quality-improvement teaching and methods to support their adoption by teams at the participating PICUs. The main outcome measures were the rate of CA-BSIs from January 2004 to September 2007 and compliance with each element of the insertion and maintenance bundles from October 2006 to September 2007.

**RESULTS:** Average CA-BSI rates were reduced by 43% across 29 PICUs (5.4 vs 3.1 CA-BSIs per 1000 central-line-days;  $P < .0001$ ). By September 2007, insertion-bundle compliance was 84% and maintenance-bundle compliance was 82%. Hierarchical regression modeling showed that the only significant predictor of an observed decrease in infection rates was the collective use of the insertion and maintenance bundles, as demonstrated by the relative rate (RR) and confidence intervals (CIs) (RR: 0.57 [95% CI: 0.45–0.74];  $P < .0001$ ). We used comparable modeling to assess the relative importance of the insertion versus maintenance bundles; the results showed that the only significant predictor of an infection-rate decrease was maintenance-bundle compliance (RR: 0.41 [95% CI: 0.20–0.85];  $P = .017$ ).

**CONCLUSIONS:** In contrast with adult ICU care, maximizing insertion-bundle compliance alone cannot help PICUs to eliminate CA-BSIs. The main drivers for additional reductions in pediatric CA-BSI rates are issues that surround daily maintenance care for central lines, as defined in our maintenance bundle. Additional research is needed to define the optimal maintenance bundle that will facilitate elimination of CA-BSIs for children. *Pediatrics* 2010;125:206–213

**AUTHORS:** Marlene R. Miller, MD, MSc,<sup>a,b,c</sup> Michael Griswold, PhD,<sup>d</sup> J. Mitchell Harris, II, PhD,<sup>c</sup> Gayane Yenokyan, MD, MPH,<sup>d</sup> W. Charles Huskins, MD,<sup>e</sup> Michele Moss, MD,<sup>f</sup> Tom B. Rice, MD,<sup>g</sup> Debra Ridling, RN, MS, CCRN,<sup>h</sup> Deborah Campbell, RNC, CCRN,<sup>i</sup> Peter Margolis, MD, PhD,<sup>j</sup> Stephen Muething, MD,<sup>j</sup> and Richard J. Brill, MD<sup>k</sup>

<sup>a</sup>Department of Pediatrics, Johns Hopkins University School of Medicine, Baltimore, Maryland; Departments of <sup>b</sup>Health Policy and Management and <sup>c</sup>Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland; <sup>d</sup>National Association of Children's Hospitals and Related Institutions, Alexandria, Virginia; <sup>e</sup>College of Medicine, Mayo Clinic, Rochester, Minnesota; <sup>f</sup>Department of Pediatrics, Arkansas Children's Hospital, Little Rock, Arkansas; <sup>g</sup>Critical Care, Children's Hospital of Wisconsin, Milwaukee, Wisconsin; <sup>h</sup>Nursing Quality and Evidence-Based Practice, Seattle Children's Hospital, Seattle, Washington; <sup>i</sup>Pediatric Critical Care, Kosair Children's Hospital, Louisville, Kentucky; <sup>j</sup>Divisions of Health Policy and Clinical Effectiveness and General Pediatrics, Center for Child Health Quality, Cincinnati, Ohio; and <sup>k</sup>Division of Critical Care Medicine, Cincinnati Children's Hospital, Cincinnati, Ohio

### KEY WORDS

bloodstream infections, children, catheter-associated bloodstream infections, nosocomial infections, pediatric intensive care unit, hospital-acquired infections

### ABBREVIATIONS

CA-BSI—catheter-associated bloodstream infection  
RR—relative rate  
CI—confidence interval  
CDC—Centers for Disease Control and Prevention  
NACHRI—National Association of Children's Hospitals and Related Institutions  
CVL—central venous line

Dr Brill's current affiliation is Department of Pediatrics, Nationwide Children's Hospital, Columbus, OH.

[www.pediatrics.org/cgi/doi/10.1542/peds.2009-1382](http://www.pediatrics.org/cgi/doi/10.1542/peds.2009-1382)

doi:10.1542/peds.2009-1382

Accepted for publication Aug 3, 2009

Address correspondence to Marlene R. Miller, MD, MSc, Johns Hopkins University, Johns Hopkins Children's Center, CMSC 1-141, 600 N Wolfe St, Baltimore, MD 21287. E-mail: mmille21@jhmi.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2010 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

Catheter-associated bloodstream infections (CA-BSIs) are a significant cause of morbidity, mortality, and added medical costs to hospitalized adult and pediatric patients.<sup>1–11</sup> According to the National Nosocomial Infection Surveillance System established by the Centers for Disease Control and Prevention (CDC), the pooled mean in 2004 among 54 PICUs was 6.6 CA-BSIs per 1000 catheter-days, higher than in many adult ICUs.<sup>12</sup> More recent estimates that included 36 PICUs have shown a pooled mean rate of 5.3 CA-BSIs per 1000 catheter-days.<sup>13</sup> Despite these data, strategies to reduce health care complications such as pediatric CA-BSIs have received relatively little attention.

CA-BSIs in adult ICUs have been nearly eliminated by applying a multifaceted intervention<sup>14–17</sup> that is now used broadly throughout the United States and the world. Whether these findings and interventions apply to children is not known.

Children who receive care in PICUs have unique, albeit not thoroughly researched, risk factors for CA-BSIs compared with adult patients. These differences include a less-clear protective effect of central lines in the jugular or subclavian veins, as opposed to the femoral veins, and the presence of underlying genetic syndromes and congenital malformations.<sup>18–20</sup> In addition, pediatric providers use patient central lines differently. For example, consistent anecdotal reports have shown providers to have a greater reliance on central lines to obtain needed blood samples or to keep the lines in place longer to ensure secure venous access in an emergency. The National Association of Children's Hospitals and Related Institutions (NACHRI) developed a quality-improvement collaborative, composed of 29 PICUs, to identify and test the impact of pediatric-

specific catheter-care practices in reducing pediatric CA-BSI rates.

## METHODS

### Study Design, Setting, Participants, and Objectives

Twenty-seven NACHRI member hospitals agreed to work collaboratively to reduce the CA-BSI rates among their 29 PICUs. The study design was a multi-institutional, interrupted time-series design that included historical control data from each of the 29 PICUs from January 2004 to August 2006. With a goal of eliminating CA-BSIs attributed to the PICU, the collaborative set interim first-year goals of a 50% decrease in CA-BSI rates by reliably using the collaborative insertion bundle for 90% of central line insertions and by reliably performing the collaborative maintenance bundle for 70% of all central line catheter-maintenance care.

As part of their participation in the collaborative, each team committed a senior PICU leader/physician champion to support and promote the unit team. The 2 to 3 additional team members for each site included quality-improvement leaders, infectious disease physicians, PICU nursing leaders, and/or infection-control professionals. The PICU teams participated in 4 face-to-face learning workshops in the first year, monthly conference calls, and monthly data collection and submission. Each PICU team, using quality-improvement methods of small tests of change, tested and implemented changes to make their care practices commensurate with the collaborative's recommended central line insertion and maintenance-care practices. From October 2006 through September 2007, the teams engaged in intensive efforts to reduce PICU CA-BSI rates by implementing the insertion and maintenance-care bundles. Each PICU team consulted with its respective in-

stitutional review board (IRB) regarding collaborative participation and received either IRB approval or a determination that the effort did not need IRB review.

### Interventions

Our study involved 2 central line care bundles, one focused on insertion practices and one focused on maintenance practices (Table 1). The insertion bundle included evidence-based procedures recommended by the CDC that have been proven to be effective in adult patients or in a single institutional PICU.<sup>16,21,22</sup> Contrary to adult CA-BSI efforts, our bundle did not discourage the use of the femoral site, because this can often be the most convenient or optimal insertion site for children, and the evidence of increased infection risk with femoral sites is not definitive for pediatric patients.<sup>12,23–25</sup> The maintenance bundle was created by using some of the pertinent CDC guidelines; however, consensus of mostly expert pediatric physicians and nurses from approximately 20 children's institutions were involved in the development of this effort because of the relative paucity of more rigorous evidence on an effective maintenance bundle.<sup>14,21</sup>

### Measures and Data

Data were collected by using insertion-bundle and maintenance-bundle compliance as the 2 process measures. Each PICU team self-monitored all central-line insertions that occurred in the PICU and submitted data on compliance with each insertion-bundle element for all of the insertions that occurred each month. Once each week, each PICU team audited all of the central lines to check for compliance with each element of the maintenance bundle. The insertion-bundle and maintenance-bundle audit data were recorded each month by each PICU and entered into the collaborative data-

**TABLE 1** Central Line Catheter–Care Bundles

Insertion bundle
Wash hands before the procedure.
For all children aged $\geq 2$ mo, use chlorhexidine gluconate to scrub the insertion site for 30 s for all areas except the groin, which should be scrubbed for 2 min. Scrubbing should be followed by 30 to 60 s of air drying.
No iodine skin prep or ointment is used at the insertion site.
Prepackage or fill the insertion cart, tray, or box including full sterile barriers
Create an insertion checklist, which empowers staff to stop a nonemergent procedure if it does not follow sterile insertion practices.
Use only polyurethane or Teflon catheters. <sup>a</sup>
Conduct insertion training for all care providers, including slides and video.
Maintenance bundle
Assess daily whether catheter is needed.
Catheter-site care
No iodine ointment.
Use a chlorhexidine gluconate scrub to sites for dressing changes (30-s scrub, 30-s air-dry).
Change gauze dressings every 2 d unless they are soiled, dampened, or loosened. <sup>a</sup>
Change clear dressings every 7 d unless they are soiled, dampened, or loosened. <sup>a</sup>
Use a prepackaged dressing-change kit or supply area
Catheter hub, cap, and tubing care
Replace administration sets, including add-on devices, no more frequently than every 72 h unless they are soiled or suspected to be infected.
Replace tubing that is used to administer blood, blood products, or lipids within 24 h of initiating infusion. <sup>a</sup>
Change caps no more often than 72 h (or according to manufacturer recommendations); however, caps should be replaced when the administration set is changed. <sup>a</sup>
The prepackaged cap-change kit, or supply area elements to be designated by the local institution.

<sup>a</sup> These procedures are according to the CDC recommendations.

base. Compliance for both bundles was assessed as all or none, meaning that each patient's insertion or maintenance event had to comply with all of the elements of the respective bundle to be considered compliant.

Our outcome measure was the monthly PICU CA-BSI rate, defined as CA-BSI cases per 1000 central-line-days. The number of CA-BSI cases and the monthly total of central-line-days per PICU were collected by trained, hospital-based, infection-control practitioners in accordance with CDC definitions.<sup>21</sup>

### Analysis and Interpretation

Monthly and quarterly data pertaining to the CA-BSI collaborative were recorded and analyzed from January 2004 through September 2007. The entire collaborative effort was subdivided into 3 time periods for the analysis. The baseline period was from January 2004 through September 2006, quarters 1 through 11. A ramp-up

period was defined as the initial 3 months of collaborative interventions from October 2006 until December 2006, when there was variation between the units regarding how many of the bundle elements were being fully implemented. A stable-effect period ranged from January 2007 through September 2007 during sustained and comprehensive CA-BSI intervention efforts by all of the PICU teams.

To examine the data and account for the clustering effect within the individual PICUs, we used hierarchical modeling including marginal generalized linear models with log-links, negative binomial distributions, and working autoregressive correlation structures. These models were fit by using generalized estimating equations with robust variance estimation.<sup>26</sup> The analytic model allowed the CA-BSI rate to change as a function of time within each period (different temporal slopes within the baseline, ramp-up, and

stable-effects periods); the log-link in the generalized linear models implied that exponentiated coefficients were interpreted as relative rates (RRs), which were relative increases or decreases above or below the last quarter's rate, similar to a relative risk. Specifically, the model estimated the RR (increase or decrease) of CA-BSIs per quarter during the baseline period, the RR of CA-BSIs comparing the ramp-up period to the last quarter of the baseline period, and the RR of CA-BSIs per quarter during the stable-effects period. An additional simplified model was examined to estimate a single CA-BSI rate during each of the 3 periods (baseline, ramp-up, and stable-effects) and to compare those rates. The use of the simplified model assuming a single CA-BSI rate during each of the 3 periods was considered acceptable given that within each period, the CA-BSI rate did not significantly change over time. Models were adjusted for geographic region, average length of stay, and bed capacity to account for any unpredicted regional variations in care and to attempt to account for any PICU-specific severity of patient illness. Nonlinear relationships were explored by using the lowest smoothing functions and splines.<sup>27,28</sup>

Hierarchical regression modeling was also used to examine individual compliance effects (either insertion or maintenance) after adjusting for covariables and the alternate compliance variable.<sup>29</sup>

Missing data in our analyses were exclusively confined to the baseline pre-collaborative period (January 2004 to September 2006) and encompassed 18.6% of baseline data. To examine the effects of the missing data, we ran 2 sensitivity analyses models by imputing data for each missing data point. All of the results were similar in estimates and statistical significance to

the all-observable analyses reported here.<sup>30–33</sup>

All analyses were conducted by using SAS 9.1 (SAS Institute, Inc, Cary, NC), and statistical process control charts, including U charts, were also used in the preliminary analyses to examine whether the observed differences were a special-cause variation or a common-cause variation.<sup>34</sup>

## RESULTS

Table 2 provides the characteristics of the 29 pediatric PICUs enrolled in our NACHRI PICU CA-BSI collaborative. Overall, most of the PICUs were mixed pediatric and cardiac PICUs, with 2 being solely pediatric cardiac ICUs. The majority of sites had level 1 trauma centers

**TABLE 2** Characteristics of the 29 Participating ICUs

Characteristic	<i>n</i>
Unit type	
PICU	9
PICU/CICU	18
CICU	2
Beds, <i>n</i>	
10–16	12
17–27	13
28–36	4
Annual PICU length of stay, mean, d	
2.7–4.5	14
4.6–6.3	12
6.4–9.6	3
Yearly total of PICU patient days, d	
2100–3600	10
3700–6200	10
6300–8700	9
ICU annual admissions in 2005	
300–899	10
900–1700	15
1800–2400	4
Institution is level 1 trauma center	
Yes	19
No	10
Institution performs solid-organ transplants	
Yes	25
No	4
Institution performs bone marrow transplants	
Yes	21
No	7
ICU performs ECMO	
Yes	26
No	3

CICU indicates cardiac ICU; ECMO, extracorporeal membrane oxygenation.

and performed solid-organ transplants, bone marrow transplants, and extracorporeal membrane oxygenation.

The average baseline CA-BSI rate for the 29 PICUs, in aggregate, was 5.4 CA-BSIs per 1000 central-line-days. After initiation of the interventions, the rate began to decrease in the ramp-up period (4.3 CA-BSIs per 1000 central-line-days) and decreased to an average stable-effect rate of 3.1 CA-BSIs per 1000 central-line-days. These results translate into a significant difference between the baseline CA-BSI rate and the steady-state stable-effect period CA-BSI rate, with a decrease from 5.4 (95% confidence interval [CI]: 4.5–6.4–3.1 and 2.4–4.0, respectively;  $P < .0001$ ).


In terms of the stated first-year goals of the NACHRI PICU CA-BSI collaborative, the 29 PICUs achieved a 43% reduction in CA-BSI rates. By the end of the first year, the collaborative came close to achieving the insertion-bundle compliance goal with 84% sustained compliance and met the maintenance-bundle compliance goal with 82% sustained compliance.

The 29 PICUs reported 324 CA-BSI events during the 12-month postinter-

vention study period (95 205 total central-line-days).

We used hierarchical cluster-analysis regression modeling and adjusted for the ICU region and PICU demographics (average length of stay and bed capacity). The only significant predictor of the observed decrease in CA-BSI rates was the collective interventions that were used in this collaborative, namely the insertion and maintenance bundles, during the stable-effect period (RR: 0.57 [95% CI: 0.45–0.74];  $P < .0001$ ). Given the downward trend in CA-BSI rates in the baseline data, a more-stringent sensitivity analysis, assuming that the baseline period CA-BSI rate was actually decreasing as opposed to being constant, was conducted to account for this trend. Nearly identical results were found (RR: 0.618 [95% CI: 0.47–0.82];  $P < .001$ ).

To begin exploring which of the bundles had the greatest influence on the observed CA-BSI rate decrease, Fig 1 shows the lowest smoothing line and 95% CIs of the aggregate data from the 29 PICUs for both precollaborative and collaborative CA-BSI rates, as well as the lowest smoothing lines and 95%



**FIGURE 1**

Data from 29 PICUs showing the rates of CA-BSIs and insertion and maintenance compliance and 95% CIs in the precollaborative and collaborative periods.

CIs of the average compliance with the insertion and maintenance bundles in the collaborative period across all 29 PICUs. It is noteworthy that the apparent wide variation in CA-BSI rates is largely a result of variation in the denominator size among the 29 PICUs, from lows of 60 central-line-days per month to highs of 500 central-line-days per month.

Additional analysis to determine the relative importance of insertion-bundle compliance compared with maintenance-bundle compliance was performed by using hierarchical cluster modeling. After adjusting for region and PICU demographics, the only significant predictor of the CA-BSI rate decrease was maintenance-bundle compliance (RR: 0.41 [95% CI: 0.20–0.85];  $P = .017$ ).

## DISCUSSION

In this PICU CA-BSI collaborative, we demonstrated that reliably implementing a pediatric-specific central-line insertion bundle and a new pediatric central-line maintenance bundle can achieve and sustain significant decreases in pediatric CA-BSI rates in the PICU setting. In contrast with adult ICU care, maximizing insertion-bundle compliance alone cannot help PICUs eliminate CA-BSIs. Instead, the main drivers for further reducing pediatric CA-BSI rates seem to be issues that surround daily maintenance care for central lines, an attribute of bedside nursing care and practice.

To our knowledge, this study is the first to clearly articulate the differential impact of insertion-related practices versus maintenance-related practices on CA-BSI rates in either adult or pediatric populations.<sup>15–17</sup> Indeed, researchers have emphasized that practices related solely to the insertion of central lines are able, in adult patients, to significantly reduce CA-BSIs.<sup>15,16</sup> In our study, the main driver to reduce CA-

BSIs was a care bundle focused on maintenance practices, not on insertion practices.

Our analyses should not be interpreted to mean that evidence-based practices for CVL insertions with children should be abandoned. It is much more likely that the 29 PICUs in this collaborative had already maximized the impact of improving central-line insertion practices in the precollaborative period. For several years now, the health care industry has been extensively focused on insertion practices, and every PICU in our collaborative had already changed many of its insertion-related practices, on the basis of these national pressures, in the precollaborative period. In addition, our data suggest that additional reductions in CA-BSI rates will more likely be achieved by focusing attention on the day-to-day handling of central lines and not by additional work on adherence to evidence-based insertion practices. The data in Fig 1 support this assertion. At the onset of this collaborative, adherence to the ideal insertion practices was already higher than 80%. In contrast, adherence to the ideal maintenance practices at the onset of this collaborative was approximately 65%.

Our study also highlights the importance of multidisciplinary teams to address CA-BSIs and the ideal nature of collaborative models for improving pediatric care. Although nurses are clearly involved in helping to ensure adherence to ideal central-line insertion practices, this nursing role is largely a “double-check” function. In contrast, all of the clinical elements in our maintenance bundle comprise direct nursing behaviors in the day-to-day care of patients. The relationship between hands-on practices of physicians with insertion and hands-on practices of nurses for the daily care of central lines epitomizes the intertwined team nature of care and the de-

pendence on both disciplines to prevent harm to patients from CA-BSIs. Furthermore, this study showcases the strength and value of collaborative multidisciplinary improvement efforts for pediatric care.

We recognize that there are several limitations to this study. First, we did not demonstrate that this intervention can sustain a reduction in CA-BSI rates over several years. Given the track record to date of this intervention across 29 ICUs and new findings regarding the importance of the maintenance bundle, we believe that it is important to disseminate this information sooner rather than later. Our intention is to focus attention on the need for more research to further identify optimal central-line maintenance-care practices. Second, the definition of CA-BSIs as set by the CDC does not take into account the number and timing of central lines. For example, a patient with 5 simultaneous central lines over a period of 2 weeks, versus a patient with 5 sequential central lines over a period of 2 weeks, versus a patient with 1 continuous central line over a period of 2 weeks are all counted as the same number of central-line-days and assumed to be at risk for a CA-BSI, according to the CDC. These definitions, however, are standardized at all institutions that report CA-BSI data to the CDC, effectively eliminating these definitional issues as a problem when comparing institutions. Third, the 29 ICUs within the collaborative are large PICUs with an extremely ill patient population. However, the interventions described in this study could be easily implemented in other facilities regardless of their size or patient acuity. The core ideas behind these interventions revolve around educating caretakers about effective practice techniques for central-line insertion and maintenance, centralizing equipment so that adherence to best practices is easy,

reliably implementing and measuring compliance with best practices, using real-time data to drive change, and team-building including nurse empowerment. All of these attributes are applicable in any ICU setting.

## CONCLUSIONS

CA-BSIs are a preventable cause of patient harm to critically ill children. Our pediatric-specific central line insertion and maintenance bundles, which combine evidence-based guidelines with human-factors principles to reduce complexity and enhance implementation, have demonstrated that CA-BSIs are preventable for children. In contrast with adult ICU care, maximizing insertion-bundle compliance alone cannot help PICUs to eliminate CA-BSIs. Instead, the main driver for additional reductions in pediatric CA-BSI rates seems to involve issues surrounding daily maintenance care for central lines, as defined by our maintenance bundle. Although reliable implementation of best practices for central-line insertion and maintenance can significantly decrease PICU CA-BSIs, additional research is still needed to determine the optimal maintenance bundle that will facilitate elimination of CA-BSIs in all children who require short- or long-term central venous access.

## ACKNOWLEDGMENTS

We thank Kris A. Schulz, MPH, currently the senior director of research and quality improvement at the American Academy of Otolaryngology—Head and Neck Surgery. Ms Schulz previously worked at NACHRI and was instrumental in developing and launching this PICU CA-BSI collaborative. Lynne Lostocco, RN, FOCUS group director at NACHRI, and Ramesh Sachdeva, MD, PhD, JD, DBA, at Children's Hospital of Wisconsin, provided guidance and support in developing this effort. Finally, we thank all of the children's hospitals and staff who created our NACHRI PICU

CA-BSI collaborative. It is their collective hard work and dedication to improving the care of children that are presented in this article.

The NACHRI PICU CA-BSI collaborative comprises 29 centers across the United States, and includes Arkansas Children's Hospital PICU: Kim White, RN, BSN, Michelle Moss, MD, Jerril Green, MD, Craig Gilliam, and Chris Wilson, RN; Arkansas Children's Hospital CICU: Ellen Kane, Adnan Bhutta, Parthak Prodham, and Rebecca Jones, RN; Children's Hospital of Los Angeles PICU: David Schmidt, MSN, CSN, CPNP, CCRN, Barry Markovitz, MD, MPH, Wilbert Mason, MD, MPH, and Sherry Cauley, RN; UC-Davis Children's Hospital PICU: Sherri Reese, RN, CIC, Joanne Natale, MD, Robert Pretzlaff, MD, Kristi Cargill, Virpal Donley, RN, and Gina Davis, RN; Children's National Medical Center PICU: Heidi Dalton, MD, Maire Soosaar, BSN, MPH, Eveena Felder, RN, Annerose Abogadie, RN, BSN, CCRN, Arden Cooper, RN, BSN, Angela Wratney, MD, MSHc, and Christine Moran, RN, BSN; Children's National Medical Center CICU: John Berger, MD, Donna Johnson, RN, and Lisa Scafdi, RN; Alfred I. DuPont Hospital for Children PICU: Barbara Jackson, RN, BSN, Caroline Boyd, MD, and Carol Briody, MT (ASCP) CIC; Children's Hospital of Illinois at Peoria PICU: Paula LaFond, RN, Adalberto Torres, MD, Kelly Skender, PNP, Lori Goffinet, RN, and Kelly Nimtz-Rusch; Kosair Children's Hospital PICU: Justine O'Flynn, RN, Vicki Montgomery, MD, Kris Bryant, MD, Kelly Zink, Stacy Flanders, and Deborah Campbell, RN, BSN, CCRN; Johns Hopkins Children's Center PICU: Ivor Berkowitz, MD, and Judy, Ascenzi, RN; Children's Hospital Boston PICU: Greg Priebe, MD, Tom Sandora, MD, MPH, Gail Potter-Bynoe, BS, CIC, and Maggie Gellar, RN; C. S. Mott Children's Hospital PICU: Matthew Niedner, MD, Julie Juno, RN, BSN, MSc, CCRN, Allison Knapp, RN, Monica We-

ber, RN, Tonie Owens, RN, MSN, Annette Scott, RN, and Jackie Shaffer-Hartman, MPH; Helen DeVos Children's Hospital PICU: Robert Fitzgerald, MD, Rachel Blanton, and Jeni Wincek, RN, MSN; Mayo Eugenio Litta Children's Hospital PICU: W. Charles Huskins, MD, MSc, Edward Seferian, MD, Karen Stewart, RN, Alissa Struble, RN, Julie Duncan, RN, and Karen Fryer, RN; Children's Hospitals and Clinics of Minnesota, St Paul and Minneapolis PICU: Stephen Kurachek, MD, Adriene Thronton, Melanie Kuelbs, RN, and Bridget Sturtevant, RN; University of Minnesota Children's Hospital at Fairview PICU: Sandy Hagstrom, RN, CPNP, Sameer Gupta, MD, and Barbara Bor; Children's Mercy Hospital, Kansas City PICU: Patricia Webster, MD, Carolyn Greene, Maria Ginger-Wiley, RN, CCRN, Nancy Brundage, RN, and Robyn Livingston, MD; Joseph M. Sanzari Children's Hospital PICU: Stephen Percy, MD, MBA, Carol Weber, RN, and Jennifer Brady, RN; Children's Hospital of New Jersey at Newark Beth Israel Medical Center PICU: Meena Kalyanaraman, MD, Pat Harmon, Cheryle Aizley, MSN, Tim Yeh, MD, Kristin Goldmacher, RN, BSN, and Marcia Summers, RN; University of New Mexico Children's Hospital PICU: Renee Donohue, Dawn Joseph, MD, Rachel Rivera, RN, Cynthia Johnson, and April Figueroa, RN; Duke University Medical Center PICU: Jon Meliones, MD, Rebecca Ellis, RN, BSN, Kristi Ryan, RN, BSN, Lisa Cooper, RN, BSN, ICP, and Kshitij Mistry, MD, MSc; Cincinnati Children's Hospital PICU: Richard Brilli, MD, Mary Jo Giaccone, RN, MSN, Sarah Golden, RN, and Terry Palmisano, RN, MSN; Cincinnati Children's Hospital CICU: Laurie Mustin, RN, CPNP, Catherine Dent, MD, Anna Sheets, RN, and Sue Ryckman, RN, CPNP; Children's Hospital Medical Center of Akron PICU: Jean Christopher, RN, MSN, CNS, Ann-Marie Brown, MSN, CPNP, CCRN, and Michel Forbes, MD; Penn State Children's Hospital PICU: Michael Dettorre, DO, Linda

Risser, RN, BSN, Amanda Politz, RN, Megan Latsha, RN, Carolyn Foerster, RN, Emily Lewellen, RN, MSN, CRNP, and Laura Bower, RN, BSN; Cook Children's Medical Center PICU: Larry Easterling, MD, Joann Sanders, MD, Leah Hekele, RN, Kimberly Williams, RN, Rosa Boyett, RN, Jan Park, RN, Cheryl Patterson, RN, MSN, and Laurie Patterson, RN; Dell

Children's Medical Center of Central Texas PICU: LeeAnn Christie, RN, Renee Higgerson, MD, and Laura Buford, RN; Inova Fairfax Hospital for Children PICU: Elizabeth Suddaby, RN, MSN, Keith Dockery, MD, Margaret Sourbeer, RN, MSN, Nelia Bruce, RN, and Kathi Huddleston; Children's Hospital and Regional Medical Center, Seattle PICU: De-

bra Ridling, RN, CCRN, Howard Jeffries, MD, MBA, Elaine Albert, MD, Audrey Aboulafia, RN, and Julie Smith, RN; and Children's Hospital of Wisconsin, Milwaukee PICU: Tom Rice, MD, Krista Colpaert, RN, Ramesh Sachdeva, MD, Rainer Gedeit, MD, Mary Bolhuis, RN, Deb Soetenga, APN, Linda Flannery, RN, Chris Lund-Holtz, RN, and Mary Rotar, RN.

## REFERENCES

- Mermel LA. Prevention of intravascular catheter-related infections. *Ann Intern Med.* 2000;132(5):391–402
- Pittet D, Tarara D, Wenzel RP. Nosocomial bloodstream infection in critically ill patients: excess length of stay, extra costs, and attributable mortality. *JAMA.* 1994; 271(20):1598–1601
- Richards MJ, Edwards JR, Culver DH, Gaynes RP; National Nosocomial Infections Surveillance System. Nosocomial infections in pediatric intensive care units in the United States. *Pediatrics.* 1999; 103(4):e39
- Urrea M, Pons M, Serra M, Latorre C, Palomeque A. Prospective incidence study of nosocomial infections in a pediatric intensive care unit. *Pediatr Infect Dis J.* 2003; 22(6):490–494
- Yogaraj JS, Elward AM, Fraser VJ. Rate, risk factors, and outcomes of nosocomial primary bloodstream infection in pediatric intensive care unit patients. *Pediatrics.* 2002; 110(3):481–485
- Grohskopf LA, Sinkowitz-Cochran RL, Garrett DO, et al. A national point-prevalence survey of pediatric intensive care unit-acquired infections in the United States. *J Pediatr.* 2002;140(4):432–438
- Odetola FO, Moler FW, Dechert RE, VanDerElzen K, Chenoweth C. Nosocomial catheter-related bloodstream infections in a pediatric intensive care unit: risk and rates associated with various intravascular technologies. *Pediatr Crit Care.* 2003;4(4): 432–436
- Wisplinghoff H, Seifert H, Tallent S, Bischoff T, Wenzel RP, Edmond MB. Nosocomial bloodstream infections in pediatric patients in United States hospitals: epidemiology, clinical features and susceptibility. *Pediatr Infect Dis J.* 2003; 22(8):686–691
- Aly H, Herson V, Duncan A, Herr J, Bender J, Patel K, El-Mohandes AAE. Is bloodstream infection preventable among premature infants? A tale of two cities. *Pediatrics.* 2005; 115(6):1513–1518
- Elward AM, Hollenbeak CS, Warren DK, Fraser VJ. Attributable cost of nosocomial primary bloodstream infection in pediatric intensive care unit patients. *Pediatrics.* 2005;115(4):868–872
- Costello J, Morrow D, Graham D, Potter-Bynoe G, Sandora TJ, Laussen PC. Systematic intervention to reduce central line-associated bloodstream infection rates in a pediatric cardiac intensive care unit. *Pediatrics.* 2008;121(5):915–923
- National nosocomial infections surveillance (NNIS) system report, data summary from January 1992 through June 2004, issued October 2004. *Am J Infect Control.* 2004;32(8): 470–485
- Edwards JR, Peterson KD, Andrus ML, et al. National Healthcare Safety Network (NHSN) Report, data summary for 2006, issued June 2007. *Am J Infect Control.* 2007;35(5): 290–301
- O'Grady NP, Alexander M, Dellinger, EP, et al. Guidelines for the prevention of intravascular catheter-related infections. *Pediatrics.* 2002;110(5):e51
- Berenholtz SM, Pronovost PJ, Lipsett PA, et al. Eliminating catheter-related bloodstream infections in the intensive care unit. *Crit Care Med.* 2004;32(10):2014–2020
- Pronovost P, Needham D, Berenholtz S, et al. An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med.* 2006;355(26):2725–2732
- Institute for Healthcare Improvement. Prevent central line-associated blood stream infections. Available at: [www.ihl.org/IHI/Programs/Campaign/Centrallineinfection.htm](http://www.ihl.org/IHI/Programs/Campaign/Centrallineinfection.htm). Accessed December 18, 2008
- Elward A, Fraser V. Risk factor for nosocomial primary bloodstream infection in pediatric intensive care unit patients: a 2-year prospective cohort study. *Infect Control Hosp Epidemiol.* 2006;27(6):553–560
- Kritchevsky SB, Braun, BI, Kusek L, et al. The impact of hospital practice on central venous catheter-associated bloodstream infection rates at the patient and unit level: a multicenter study. *Am J Med Qual.* 2008; 23(1):24–38
- Almuneef MA, Memish ZA, Balkhy HH, Hijazi O, Cunningham G, Francis C. Rate, risk factors and outcomes of catheter-related bloodstream infection in a paediatric intensive care unit in Saudi Arabia. *J Hosp Infect.* 2006;62(2):207–213
- National Healthcare Safety Network, Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. Available at: [www.cdc.gov/ncidod/dhqp/pdf/nnis/NosInfDefinitions.pdf](http://www.cdc.gov/ncidod/dhqp/pdf/nnis/NosInfDefinitions.pdf). Accessed December 18, 2008
- McKee C, Berkowitz I, Bradley K, et al. Reduction of catheter-related bloodstream infections in pediatric patients: experimentation and reality. *Pediatr Crit Care Med.* 2008; 9(1):40–46
- Venkataraman ST, Thompson AE, Orr RA. Femoral vascular catheterization in critically ill infants and children. *Clin Pediatr (Phila).* 1997;36(6):311–319
- Stenzel JP, Green TP, Furhman BP, Carlson PE, Marchessault RP. Percutaneous femoral venous catheterizations: a prospective study of complications. *J Pediatr.* 1989; 114(3):411–415
- Goldstein Am, Weber JM, Sheridan RL. Femoral venous access is safe in burned children: an analysis of 224 catheters. *J Pediatr.* 1997;130(3):442–446
- Diggle PJ, Heagerty P, Liang KY, Zeger SL. *Analysis of Longitudinal Data.* Oxford, United Kingdom: Oxford University Press; 2002
- Durrleman S, Simon, R. Flexible regression models with cubic splines. *Stat Med.* 1989; 8(5):551–561
- Hastie TJ, Tibshirani RJ. *Generalized Additive Models.* Boca Raton, FL: Chapman and Hall/CRC; 1990
- Wang PC. Adding a variable in generalized

- linear models. *Technometrics*. 1985;27(3): 273–276
30. Little RJA, Rubin DB. *1987 Statistical Analysis With Missing Data*. 2nd ed. New York, NY: John Wiley and Sons; 2002
31. Rubin DB. *1987 Multiple Imputation for Non-response in Surveys*. New York, NY: John Wiley and Sons; 2004
32. Rubin DB. Multiple imputation after 18+ years (with discussion). *J Am Stat Assoc*. 1996;91(434):473–489
33. Schafer JL. *Analysis of Incomplete Multivariate Data*. London, United Kingdom: Chapman & Hall; 1997
34. Benneyan JC, Lloyd RC, Plsek PE. Statistical process control as a tool for research and healthcare improvement. *Qual Saf Health Care*. 2003;12(6):458–464

**Decreasing PICU Catheter-Associated Bloodstream Infections: NACHRI's  
Quality Transformation Efforts**

Marlene R. Miller, Michael Griswold, J. Mitchell Harris, II, Gayane Yenokyan, W.  
Charles Huskins, Michele Moss, Tom B. Rice, Debra Ridling, Deborah Campbell,  
Peter Margolis, Stephen Muething and Richard J. Brill

*Pediatrics* published online Jan 11, 2010;

DOI: 10.1542/peds.2009-1382

**Updated Information  
& Services**

including high-resolution figures, can be found at:  
<http://www.pediatrics.org>

**Permissions & Licensing**

Information about reproducing this article in parts (figures,  
tables) or in its entirety can be found online at:  
<http://www.pediatrics.org/misc/Permissions.shtml>

**Reprints**

Information about ordering reprints can be found online:  
<http://www.pediatrics.org/misc/reprints.shtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

